

REVIEW PAPER

# At the border: the plasma membrane–cell wall continuum

Zengyu Liu<sup>1</sup>, Staffan Persson<sup>1,2</sup> and Clara Sánchez-Rodríguez<sup>1,3,\*</sup>

<sup>1</sup> Max-Planck Institute for Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam, Germany

<sup>2</sup> ARC Centre of Excellence in Plant Cell Walls, School of Botany, University of Melbourne, Parkville 3010, Victoria, Australia

<sup>3</sup> Department of Biology, ETH Zurich, 8092 Zurich, Switzerland

\* To whom correspondence should be addressed. E-mail: [rodriguez@mpimp-golm.mpg.de](mailto:rodriguez@mpimp-golm.mpg.de)

Received 10 October 2014; Revised 23 December 2014; Accepted 2 January 2015

## Abstract

Plant cells rely on their cell walls for directed growth and environmental adaptation. Synthesis and remodelling of the cell walls are membrane-related processes. During cell growth and exposure to external stimuli, there is a constant exchange of lipids, proteins, and other cell wall components between the cytosol and the plasma membrane/apoplast. This exchange of material and the localization of cell wall proteins at certain spots in the plasma membrane seem to rely on a particular membrane composition. In addition, sensors at the plasma membrane detect changes in the cell wall architecture, and activate cytoplasmic signalling schemes and ultimately cell wall remodelling. The apoplastic polysaccharide matrix is, on the other hand, crucial for preventing proteins diffusing uncontrollably in the membrane. Therefore, the cell wall–plasma membrane link is essential for plant development and responses to external stimuli. This review focuses on the relationship between the cell wall and plasma membrane, and its importance for plant tissue organization.


**Key words:** Anchor proteins, cell wall, plasma membrane, signalling, vesicle trafficking.

## Introduction

Cell walls are mechanistically strong and dynamic structures that encase plant cells. The cell walls protect the cells against environmental stresses and play key roles in directed plant growth and cell differentiation (Somerville *et al.*, 2004; Tsukaya, 2006; Ivakov and Persson, 2013), and are mainly composed of polysaccharides, proteins, and polyphenols (Boerjan *et al.*, 2003; Rose, 2003; Jamet *et al.*, 2006). All growing plant cells are surrounded by primary cell walls, which in dicot species comprise cellulose, hemicelluloses, pectins, and proteins, such as RGD (Arg-Gly-Asp tripeptide) recognizing proteins and glycosylphosphatidylinositol (GPI) anchored proteins (Canut *et al.*, 1998; Cosgrove, 2005; Gillmor *et al.*, 2005). Additional secondary cell wall layers are produced after cessation of cell expansion in some specialized cells. These walls typically provide mechanical support to the plant body through their ordered cellulose fibrils and lignin content (Boerjan *et al.*, 2003).

With the exception of cellulose, cell wall polysaccharides and proteins are synthesized in the endoplasmic reticulum and

Golgi apparatus, and secreted to the apoplast via vesicle trafficking. Cellulose is synthesized at the plasma membrane by transmembrane cellulose synthase (CesA) complexes (CSCs), which are assembled in either the endoplasmic reticulum or Golgi, and subsequently delivered to the plasma membrane (Lerouxel *et al.*, 2006; Geisler *et al.*, 2008). Once at the plasma membrane, the CSCs synthesize unbranched glucan chains that assemble into para-crystalline cellulose microfibrils (Morgan *et al.*, 2013; Fig. 1). The microfibrils become fixed in the cell wall and the catalytic capacity of the CSCs has therefore been proposed to exert a force that moves the complex forward inside the plasma membrane during cellulose synthesis. This hypothesis is supported by recent structural estimations of cellulose synthase from *Rhodobacter* (Morgan *et al.*, 2013; Sethaphong *et al.*, 2013). Thus, the motility of the CSCs should reflect the rate of cellulose production and might be dependent on the composition of the plasma membrane. The CSCs have been shown to have an average lifetime



**Fig. 1.** The plasma membrane–cell wall anchors. Cell walls are mainly composed of cellulose microfibrils and matrix polysaccharides. Each cellulose microfibril is formed by glucan chains assembled and translocated by a CSC, which moves in the plasma membrane guided by cortical microtubules through CSI1. The cell wall and plasma membrane are physically linked by plasma membrane proteins, characterized by particular domains, which anchor apoplastic polysaccharides. Thus GPI is necessary for anchoring certain proteins to the plasma membrane, such as COBRA and fasciclins. Other proteins, including integrin-like and receptor like kinases, such as WAKs, are attached to the plasma membrane through transmembrane domains. The ability of these proteins to bind to the cell wall relies on the presence of cellulose-binding domains (in COBRA) or arabinogalactan moieties which covalently attach to matrix polysaccharides (as in the case of arabinogalactan proteins and FLAs). Integrin-like proteins can bind to the RGD domain of apoplastic proteins, which may bind cell wall polysaccharides. However, many questions are still open regarding these interactions, since no RGD-containing protein has been identified in plants.

at the plasma membrane of 10–20 min (Paredes, *et al.*, 2006; Gutierrez *et al.*, 2009; Sampathkumar *et al.*, 2013), suggesting that, after they have fulfilled their function, they are internalized via endocytosis. This process also removes lipids, other proteins, and probably cell wall constituents from the plasma membrane and apoplast. Hence, cell wall architecture and remodelling depends on efficient exchange of material between the cell's interior and the plasma membrane/apoplastic space (Drakakaki and Dandekar, 2013; Sánchez-Rodríguez and Persson, 2014).

While being synthesized and remodelled, the cell wall is physically anchored to the plasma membrane via the cellulose fibrils and plasma membrane proteins that interact with cell wall components (Baluška *et al.*, 2003). This anchoring is essential for proteins to retain their localization at the plasma membrane (e.g. PIN2 and other proteins: Feraru *et al.*, 2011; Martinière and Runions, 2013), for cell wall synthesis and remodelling (e.g. during Casparian strip formation: Alassimone *et al.*, 2010; Roppolo and Geldner, 2012), and for signal transduction (Wolf *et al.*, 2012a).

Taken together, the cell wall–plasma membrane interactions comprise four main areas: the plasma membrane as a gating point for cell wall components; the plasma membrane as a plane for CSC movement; the plasma membrane–cell wall cross-links; and cell wall signalling via plasma membrane-located signalling transmitters. Considering the vast number of reviews dealing with the first, second and fourth of these areas (Wolf *et al.*, 2012a; Bashline *et al.*, 2014; McFarlane *et al.*, 2014; Sánchez-Rodríguez and Persson, 2014), we only briefly review and provide recent updates of these, and instead focus on cross-linking between the plasma membrane and cell wall in plant cells.

## The plasma membrane: the ‘customs’ of the plant cell

Many biological components are made in one location but needed in another. This certainly holds true for the majority of cell wall components, which are synthesized inside the cell, but assembled at the apoplast. Therefore, efficient membrane trafficking is important for cell wall formation and remodelling (for recent reviews see Bashline *et al.*, 2014; McFarlane *et al.*, 2014; Sánchez-Rodríguez and Persson, 2014).

Soluble cell wall polysaccharides; i.e. pectins and hemicelluloses, have been found in post-Golgi vesicles together with enzymes involved in their synthesis (Young *et al.*, 2008; Driouch *et al.*, 2012; Atmodjo *et al.*, 2013). These observations support the hypothesis that soluble cell wall polysaccharides are synthesized inside the cells, and that they are translocated to the apoplast by secretion. During exocytosis, vesicles derived from the trans-Golgi network (TGN) traffic through the cytosol via actin filaments and bundles, and fuse to the plasma membrane, which leads to the release of their content into the existing cell wall (Drakakaki and Dandekar, 2013; Sánchez-Rodríguez and Persson, 2014). Recently, an ECH/YIP complex-dependent pathway was found to promote the formation of soluble polysaccharide-containing

vesicles at the TGN (Gendre *et al.*, 2013). Accordingly, lesions in members of the complex led to alterations in cell elongation, seed coat mucilage deposition, and anther and pollen development (Gendre *et al.*, 2011; Fan *et al.*, 2014). After vesicle shuttling through the cytosol, the vesicles need to be targeted and tethered to the plasma membrane before membrane fusion. The SEC8 and EXO70A1 subunits of the tethering exocyst complex, and their interactor ROH1, have been reported to influence pectin deposition in the seed coat (Kulich *et al.*, 2010), indicating that the exocyst secretes soluble polysaccharides. Other cell wall components, e.g. lignin precursors, can be translocated from the cytosol to the apoplast by plasma membrane-localized ABC transporters (McFarlane *et al.*, 2010; Kang *et al.*, 2011; Alejandro *et al.*, 2012). In contrast, cellulose is synthesized at the plasma membrane. Therefore, efficient transport of the CSCs to and from the plasma membrane is important for cellulose synthesis (Sampathkumar *et al.*, 2013; McFarlane *et al.*, 2014).

Morphometric studies have estimated that around 75% of the total membrane incorporated into the plasma membrane of an expanding cell, or cell plate during cytokinesis, is recycled by endocytosis (Samuels and Bisalputra, 1990; Ketelaar *et al.*, 2008). Hence, a finely tuned balance of exo- and endocytosis appears to be important for proper cell wall production. This hypothesis is supported by recent data, which indicate that clathrin-mediated endocytosis (CME) is required for cell wall biogenesis, in particular for cellulose synthesis. Various CME proteins have been reported to be essential for plant cell division and cell elongation based on their localization and on cellulose deficiencies in corresponding mutants (Kang *et al.*, 2003; Van Damme *et al.*, 2006; Collings *et al.*, 2008; Frigerio, 2010; Xiong *et al.*, 2010; Bashline *et al.*, 2013; Kim *et al.*, 2013; Miart *et al.*, 2014). However, the mechanism by which the CME subunits internalize CSCs is still being elucidated. Interestingly, endocytosis of the GPI-anchored proteins, involved in cell adhesion, has been reported in animals (Lakhan *et al.*, 2009). Related plant cell wall protein might, therefore, also be internalized from the plasma membrane; however, this remains to be proven. In addition, although the data are inconclusive, internalization of pectins and hemicelluloses by CME is possible (Baluška *et al.*, 2002; Baluška *et al.*, 2005).

## Movement of CSCs in the plasma membrane

Recent observations indicate that the CSCs are preferentially inserted into the plasma membrane adjacent to cortical microtubules, but how this is achieved is not yet known (Gutierrez *et al.*, 2009). It is plausible that the membrane composition is important for this, as it seems to play a key role in exo- and endocytosis (Bessueille *et al.*, 2009; Žárský *et al.*, 2009; Zhang *et al.*, 2014). Proteomic analysis of detergent-resistant membrane (DRM) fractions extracted from various plant species revealed that the plasma membrane domains contain many different plasma membrane proteins (Morel *et al.*, 2006; Lefebvre *et al.*, 2007; Bessueille *et al.*, 2009), including CesAs (Bessueille

*et al.*, 2009). However, the CesA proteins were not identified in other proteomic studies of DRMs (Borner *et al.*, 2005; Morel *et al.*, 2006; Lefebvre *et al.*, 2007), probably due to the different plant species and different extraction methods used. Indeed, the large size of the CSCs makes them difficult to isolate and identify by mass spectroscopy, even in enriched total membrane fractions, where they are abundant (Alexandersson *et al.*, 2004). Therefore, although still controversial, it is hypothesized that the CSCs are delivered to DRM microdomains.

Lipid analysis revealed that DRMs are enriched in sterols and sphingolipids (Borner *et al.*, 2005; Lefebvre *et al.*, 2007). Sterols are one of the most important regulators of plasma membrane microdomain maintenance (Zauber *et al.*, 2014). Perturbations in sterol biosynthesis, either via inhibitors or mutations, lead to reduced cellulose content (Schrack *et al.*, 2004), indicating that sterols promote cellulose synthesis (reviewed in Schrack *et al.*, 2012). However, sterols are also implicated in various metabolic pathways, such as the biosynthesis of sitosterol- $\beta$ -glucoside (Peng *et al.*, 2002) and the phytohormone brassinosteroid, which also contribute to cellulose synthesis (Xie *et al.*, 2011; Wolf *et al.*, 2012b). Therefore, the cellulose deficiencies observed in sterol-impaired plants could be a consequence of such metabolic pathways rather than membrane composition defects. More biochemical and cell biology data are needed to clarify the direct influence of plasma membrane sterols on cellulose synthesis.

The rather imposing size of the CSCs (the membrane-spanning region has a diameter of about 30 nm, and the cytosolic domain at least twice this diameter) triggers interesting questions about how the movement of the complex in the plasma membrane affects the membrane structure and local composition. Membrane elasticity was included as a key factor in a biophysical model of CSC movement in the plasma membrane (Diotallevi and Mulder, 2007). In addition, recent computational simulations showed that the plasma membrane has an important function in the initiation of glucan assembly (Haigler *et al.*, 2014), maintaining a non-crystalline group of chains at the base of the fibril. However, experimental data on how the membrane composition affects CSC trafficking and cellulose synthesis are missing.

The CSC has been shown to be associated with other plasma membrane proteins to fulfil its function. KORRIGAN1 (KOR1), a membrane-localized endo- $\beta$ -1,4-glucanase, interacts with the CSC, influences the synthesis of cellulose, and increases the amount of non-crystalline cellulose (Nicol *et al.*, 1998; Lane *et al.*, 2001; Takahashi *et al.*, 2009; Lei *et al.*, 2014; Vain *et al.*, 2014). Cellulose Synthase Interactor 1 (CSII) is necessary for the alignment of CSC trajectories and cortical microtubules, and might bind to plasma membrane phospholipids through a C2-domain at its C-terminus (Bringmann *et al.*, 2012; Li *et al.*, 2012; Mei *et al.*, 2012). Considering their plasma membrane localization, both KOR1 and CSII might influence the CSC–membrane connexion.

## The plasma membrane–cell wall continuum

In living plant cells, the plasma membrane and cell wall are in close contact, largely due to the immense turgor pressure

(similar to that of a car tyre in growing cells) from within the cell that pushes the plasma membrane against the cell wall (Proseus and Boyer, 2005). Physical links between the plasma membrane and the cell wall are well established, as visualized by membranous structures from the plasma membrane that remain firmly anchored to the cell wall under stress generated by partial cell plasmolysis. These plasma membrane–cell wall connections are called Hechtian strands (Oparka, 1994). The Hechtian strands are mainly located at the plasmodesmata, but are also found in other plasma membrane regions (Pont-Lezica *et al.*, 1993). Although Hecht observed these strands more than 100 years ago (Hecht, 1912), the components involved in their generation and maintenance are still largely unknown.

## Cellulose microfibrils

Various data support the role of cellulose fibres in plasma membrane–cell wall adhesion (reviewed in Cvrčková, 2013; Martinière and Runions, 2013). Interestingly, cellulose has recently been shown to restrict plasma membrane protein diffusion, which may explain the slow diffusion of plant plasma membrane proteins compared to those in animal cells that lack cell walls (Owen *et al.*, 2009). These observations were obtained either by plasmolysis or by inhibiting cellulose synthesis, during which plasma membrane protein diffusion rates significantly increased (Ferarú *et al.*, 2011; Martinière *et al.*, 2011; Martinière *et al.*, 2012). Interestingly, this effect relied on the extracellular domain of the tested proteins, which included AtFH1, GFP-GPI, and PINs (Ferarú *et al.*, 2011; Martinière *et al.*, 2011; Martinière *et al.*, 2012). These data support a model in which diffusion of proteins in the plasma membrane may be regulated by interactions between their extracellular domains and cellulose (Martinière *et al.*, 2011; Martinière and Runions, 2013; Fig. 1). However, among the tested proteins above, only AtFH1 has an extracellular domain (the expansin-like domain) which may interact with cell wall polysaccharides. Therefore, the suggested interaction between cell wall and extracellular domains is not necessarily specific and the reduced lateral diffusion reported of the proteins in the presence of an intact cell wall can be explained by steric interference. Moreover, mutations in CesA subunits block the apical-basal localization of PINs in primary root cells. This suggests that the cell wall not only restricts the diffusion of plasma membrane proteins, but also supports specific localization of proteins (Ferarú *et al.*, 2011). This is in agreement with observations made with Casparian Strip Protein 1 (CASP1), which is involved in the formation of the Casparian strips in root endodermis. Indeed, CASP1 showed lateral diffusion at the lateral plasma membranes, but became exceptionally slow in the Casparian strip domain. In addition, the strong linkage of CASP1 to the extracellular matrix can only be disrupted with strong detergents. These data suggest that CASP1 might be physically linked to the cell wall via its extracellular domains, and may partially explain CASP1-specific localization (Alassimone *et al.*, 2010; Roppolo and Geldner, 2012).



Under field emission scanning electron microscopy, cellulose fibre-like structures were observed at the ends of Hechtian strands in *Tradescantia virginiana* leaf epidermal cells. The same work showed that treatment of cells with cellulase resulted in a loss of the connecting fibres (Lang *et al.*, 2004). Therefore, cellulose microfibrils were proposed to provide a physical link between the cell wall and plasma membrane. However, more recent work showed that pre-treatment of etiolated hypocotyl cells with cellulose synthesis inhibitors did not block the formation of Hechtian strands (DeBolt *et al.*, 2007). As only qualitative measurements have been done it is difficult to conclude whether the cellulose fibres actively anchor the Hechtian strands to the cell walls, or if this is achieved through other components. Interestingly, certain groups of proteins have been suggested to participate in plasma membrane–cell wall adhesion in plants, including RGD-recognizing proteins (Canut *et al.*, 1998) and GPI-anchored proteins.

### The RGD-binding proteins

RGD-dependent linkages provide a basis for cell adhesion in both plants and animals. In animals, extracellular adhesive glycoproteins typically contain RGD sequences that are recognized by plasma membrane receptors called integrins. The RGD–integrin interaction is essential for efficient cell adhesion (Alberts *et al.*, 2002). The presence of an RGD-dependent recognition mechanism and its role in cell adhesion in plants was confirmed by the application of synthetic RGD peptides to soybean suspension cultures. The exogenous RGD was here assumed to compete with the endogenous RGD-containing proteins, and this competition led to growth alterations, which were not observed when cell cultures were treated with RGE or DGR substituted proteins (Schindler *et al.*, 1989). Similar results were reported in onion and *Arabidopsis* cells, where the application of synthetic RGD peptides inhibited Hechtian strand formation (Canut *et al.*, 1998). Moreover, by immunoblotting extracts of soybean cells or protoplasts, Schindler *et al.* (1989) identified a putative plant RGD receptor that was recognized by antibodies raised against a human integrin. The RGD peptide was also found to promote the cell wall regeneration of *Nicotiana tabacum* cv. BY-2 protoplasts (Zaban *et al.*, 2013). These data indicate that plants, similarly to animals, have RGD-recognizing proteins which might help in anchoring the cell wall to the plasma membrane.

In animals, the recognition of RGD sequences by integrins activates inter- and intracellular signal transduction (Baluška *et al.*, 2003). Although plants do not have canonical integrin proteins, integrin-like proteins have been identified (Lü *et al.*, 2007a). One such example is AT14A, which contains a domain that shares high sequence similarity with animal integrins (Nagpal and Quatrano, 1999). Mutations in this protein led to a reduction in cell adhesion and an alteration in cell shape and cell wall thickness in *Arabidopsis* suspension cultures (Lü *et al.*, 2012). The Non-race-specific Disease Resistance 1 (NDR1) protein was also identified as an *Arabidopsis* integrin-like protein due to high structural similarity of its core domain to an animal integrin subunit

(Knepper *et al.*, 2011a). NDR1 contains an apoplastic RGD-like recognition motif, named NGD, and a transmembrane domain which promotes plasma membrane localization. In addition, NDR1 contains a GPI anchor (Coppinger *et al.*, 2004), which is present in many other proteins involved in plasma membrane–cell wall connections as discussed below. The receptor kinase LecRK-I.9 is a lectin-like protein with an RGD-binding motif (Bouwmeester *et al.*, 2011). Application of the recombinant LecRK-I.9 RGD-binding motif to *Arabidopsis* hypocotyls reduced the plasma membrane–cell wall contact sites, possibly by competing with the interaction between the endogenous LecRK-I.9 and RGD-binding protein (Gouget *et al.*, 2006). Subsequently, an alteration in plasma membrane–cell wall linkage was also observed in mutants altered in the receptor LecRK-I.9 (Bouwmeester *et al.*, 2011). While the plant RGD-containing protein that is recognized by LecRK-I.9 has not been identified, the RGD-binding motif of the lectin-like protein responded to a *Phytophthora infestans* effector (Senchou *et al.*, 2004). Therefore, it is plausible that RGD peptide-mediated plasma membrane–cell wall adhesion also occurs in plants (Fig. 1).

### The GPI-anchored proteins

GPI can be regarded as a surface anchor for apoplastic proteins (Fig. 1). These proteins are consequently named as GPI-anchored proteins (GAPs). The GPI anchors usually attach GAPs to microdomains in the plasma membrane (Peskan *et al.*, 2000; Bessueille *et al.*, 2009). Phospholipases (PLCs) can cleave the GPI anchor and release GAPs into the cell wall (Griffith and Ryan, 1999). The anchoring of the GAPs to the plasma membrane means that they can establish bonds to cell wall components, and may thus be regarded as promising candidates for plasma membrane–cell wall interactions. *Arabidopsis* has more than 200 putative GAPs, and 41 of these have been confirmed as targets for PLC-dependent release in callus or pollen tubes (Borner *et al.*, 2002; Borner *et al.*, 2003). The GPI-anchored proteins are essential for cell wall architecture, since mutations in GPI synthesis genes induced severe cell wall defects and growth aberrant phenotypes (Lalanne *et al.*, 2004; Gillmor *et al.*, 2005).

GAPs belong to different protein families. One of these families is represented by the *Arabidopsis* protein COBRA and its homologues found in different plants. COBRA and COBRA-like proteins have been reported to be important for anisotropic cell expansion and for the maintenance of cell wall integrity (Roudier *et al.*, 2005; Dai *et al.*, 2011; Cao *et al.*, 2012). The members of this family typically contain an N-terminal carbohydrate-binding module that can bind to crystallized cellulose *in vitro* (Liu *et al.*, 2013; Sorek *et al.*, 2014). This led the authors to hypothesize that COBRA modulates cellulose assembly, both in rice and *Arabidopsis*, by interacting with cellulose chains, which could affect microfibril crystallinity (Fig. 1). Interestingly, COBRA shows a localization pattern reminiscent of cortical microtubules in growing *Arabidopsis* cells (Roudier *et al.*, 2005). As CSCs track along cortical microtubules, these data support a role for COBRA in the assembly of cellulose fibres and their connection to the cytoskeleton. While the biochemical functions

of COBRA remain obscure, the COBRA-related proteins may thus be important factors in maintaining a plasma membrane–cell wall connection. Additional GAPs that regulate cell wall integrity exist, such as SKU5 (Sedbrook *et al.*, 2002), PMR5, and PMR6 (powdery mildew resistant gene 5 and 6: Vogel *et al.*, 2002; Vogel *et al.*, 2004), and LORELEI (Capron *et al.*, 2008; Tsukamoto *et al.*, 2010). However, the exact function of these proteins has not been clarified, and more studies are therefore required to explore their role as possible plasma membrane–cell wall anchors.

#### Arabinogalactan proteins

Arabinogalactan proteins (AGPs) are apoplastic proteins that are decorated with a glycan moiety of arabinose and galactose sugars, which accounts for more than 90% of the total mass and is suggested to be essential for AGPs function. In addition, the AGPs are typically anchored to the plasma membrane by GPI at their C-terminus (Seifert and Roberts, 2007). AGPs were shown to be covalently attached to cell wall matrix polysaccharides, i.e. to hemicellulose and pectins, by their arabinogalactan decorations (Tan *et al.*, 2013; Fig. 1). Recently, AGP31 was revealed to interact with pectins through its PAC (PRP–AGP-containing Cys) domain, which is not decorated with arabinogalactans (Hijazi *et al.*, 2014). These results indicate that the AGPs have diverse ways of interacting with different cell wall components. Many AGP mutants display defects in plant growth or cell expansion (Ellis *et al.*, 2010). For instance, a null mutation in the *Arabidopsis* classical lysine-rich AGP, AtAGP17, induced plant developmental defects during cell division and expansion (Yang *et al.*, 2007). Overexpression of its homologue, AtAGP18, resulted in bushy plants (Zhang *et al.*, 2011). In addition, AtAGP30 is involved in root regeneration and seed germination (van Hengel and Roberts, 2003). However, although the AGPs are good candidates as plasma membrane–cell wall anchors, it is unclear whether the growth-related defects seen in AGP mutants are consequences of altered plasma membrane–cell wall connections or cell wall integrity defects.

#### Fasciclin-like arabinogalactan proteins

Fasciclin-like arabinogalactan proteins (FLAs) are another class of GAPs. In *Arabidopsis*, there are 21 predicted FLAs falling into four groups (A to D), of which group A and C are GPI anchored (Johnson *et al.*, 2003). In addition, each FLA contains a fasciclin domain and one or two arabinogalactan domains (Johnson *et al.*, 2003; Fig. 1). The fasciclin domain is an extracellular module of about 140 amino acid residues with low sequence similarity, but with two highly conserved regions of ~10 amino acids each. The fasciclin domain is suggested to be an ancient cell adhesion domain common to a broad spectrum of organisms (Kawamoto *et al.*, 1998), and there are well studied cell adhesion proteins in, for example, fruitfly and human (Snow *et al.*, 1989; Elkins *et al.*, 1990; Goodman *et al.*, 1997). Based on their structure, the plant FLAs are also thought to be involved in plant plasma membrane–cell wall adhesion; however, this has not been widely explored. Interestingly, the *Arabidopsis fla4* mutant was reported to have increased plasma membrane–cell wall interface spacing, but there was not precise quantification (Shi *et al.*, 2003). Nevertheless, indirect data

support a role for plant FLAs in cell wall integrity, especially during cellulose deposition. For example, FLA4 is necessary for proper cell expansion, and for cellulose and pectin synthesis in the seeds coat (Harpaz-Saad *et al.*, 2011; Griffiths *et al.*, 2014). In addition, many *fla* mutants, in both *Arabidopsis* and cotton, showed reduced cellulose content (Li *et al.*, 2010; MacMillan *et al.*, 2010; Johnson *et al.*, 2011; Huang *et al.*, 2013).

#### Plasma membrane receptors

Another group of plasma membrane proteins that has been suggested to be plasma membrane–cell wall anchors are receptor-like kinases (RLKs) with extracellular domains that can interact with cell wall components, such as the Wall-Associated Kinases (WAKs; Kohorn and Kohorn, 2012) and members of the *Catharanthus roseus* RLK (CrRLK) family (including FERONIA, THESEUS, HERK, and ANXUR: Cheung and Wu, 2011). The extracellular domain of WAKs has been shown to bind *in vitro* to native pectin of the cell wall (Decreux and Messiaen, 2005) and, therefore, might take part in the plasma membrane–cell wall connection by covalently binding to apoplastic pectins (Wagner and Kohorn, 2001; Wolf *et al.*, 2012a). Members of the CrRLK family contain extracellular malectin-like (ML) domains with high sequence similarity to the carbohydrate-binding protein, malectin, from *Xenopus laevis* (Lindner *et al.*, 2012). Similarly, CrRLKs may interact with cell wall components through their extracellular ML domain. Indeed, a point mutation in the ML domain of THESEUS restored the growth deficiency of cellulose mutants (Hématy *et al.*, 2007). Both the WAK and CrRLK families are essential for correct cell wall growth and cell expansion (Hématy *et al.*, 2007; Fujikura *et al.*, 2014; Haruta *et al.*, 2014), and they have been previously reviewed by Cheung and Wu (2011) and Kohorn and Kohorn (2012). However, plasma membrane–cell wall linking defects in *wak* and *crrlk* mutants have not been identified.

### Cell sensing through plasma membrane–cell wall anchors

The cell wall needs to adapt its structure during cell growth, and to respond to both internal and external stresses. Therefore, physical perturbations of the cell wall must be communicated to the cell via the plasma membrane. Wolf *et al.* (2012a) recently reviewed plasma membrane receptors that might transduce cell wall-related signals. We therefore only briefly discuss RLKs in cell wall sensing, and elaborate on the putative function of other plasma membrane–cell wall anchors as sensors of cell wall signals, and their influence on plant resistance and cell morphology.

Plasma membrane–cell wall junctions may function as barriers that protect the plant cell against pathogen attacks (Mellersh and Heath, 2001). In line with this hypothesis, pre-treatment of plants with synthetic RGD peptides, which blocked plasma membrane–cell wall adhesion, increased pathogen penetration, especially of biotrophic pathogens (Mellersh and Heath, 2001). Not surprisingly, one mechanism

of biotroph pathogenicity is the disruption of plasma membrane–cell wall adhesion. Thus, the *Phytophthora infestans* RXLR-dEER effector, IPI-O, is an RGD-containing protein whose overexpression in *Arabidopsis* reduced plasma membrane–cell wall interaction (Bouwmeester *et al.*, 2011). The IPI-O is recognized by the integrin-like LecRK-I.9, and competes with the endogenous RGD-containing proteins that are made in the plant. This competition led to the destabilization of the plasma membrane–cell wall connection, benefiting progression of the pathogen. Similarly, plants mutated in LecRK-I.9 are more susceptible to *P. infestans* than wild-type plants (Bouwmeester *et al.*, 2011). In addition, mutations in the integrin-like protein NDR1 also increased plant susceptibility to pathogens (Knepper *et al.*, 2011b). Besides their function in cell expansion, WAKs have also been involved in plant immune responses (Rosli *et al.*, 2013) by recognizing pectin fragments [oligogalacturonides (OGAs)] derived from pathogen digestion of the cell wall (Brutus *et al.* 2010; Kohorn *et al.*, 2014). Interestingly, WAKs seem to distinguish the pectin native polymers from the OGAs, differentiating expansion- from stress-related signals (Kohorn *et al.*, 2014).

Cell wall mechanical sensing is another process which probably depends on the plasma membrane–cell wall interaction. A recent study showed that the plasma membrane receptor FERONIA is important for the  $\text{Ca}^{2+}$ -dependent mechanical response pathway (Shih *et al.*, 2014). Complementary work has recently shown that FERONIA activation by the secreted peptide RALF (rapid alkalization factor) inhibits plasma membrane proton activity, which induces extracellular alkalization and therefore inhibits cell expansion (Haruta *et al.*, 2014). It would be interesting to know whether FERONIA senses the cell wall components directly during the mechanical stress or senses other signals induced by the mechanical stress (like RALF and  $\text{Ca}^{2+}$ ), and subsequently alters cell growth.

The plasma membrane–cell wall connection might also be important for responses to other external stimuli, such as osmotic stress. One of the clearest outputs of osmotic stress is the production of abscisic acid (ABA) (Xiong *et al.*, 2001; Xiong *et al.*, 2002). Application of synthetic RGD peptides decreased accumulation of ABA after osmotic stress in both *Arabidopsis* and maize suspension cultures (Lü *et al.*, 2007a; Lü *et al.*, 2007b). Interestingly, a similar study using protoplasts revealed no impairment in ABA levels, corroborating the function of the plasma membrane–cell wall interaction in the osmotic stress response. Furthermore, the expression of some AGPs and FLAs was reported to respond to hormones: ATAGP30, FLA1, FLA2, and FLA8 to ABA; and ATAGP31 to jasmonic acid (van Hengel and Roberts, 2003; Liu and Mehdy, 2007). Additional studies indicate that FLAs participate in stress signalling pathways and might regulate cell wall biosynthesis. Thus, cell swelling phenotypes observed in *fla4/salt overly sensitive5* (*sos5*) mutants under elevated salt and high sucrose conditions could be rescued by the addition of external ABA (Seifert *et al.*, 2014) and ethylene biosynthesis inhibitor (Xu *et al.*, 2008), respectively. Moreover, FLA4/SOS5 was found to be genetically redundant with the FEI1/FEI2 receptor-like kinases, based on their non-additive triple

mutant phenotype (Xu *et al.*, 2008). The double mutant *fei1 fei2* displays alterations in anisotropic cell expansion and cell wall synthesis, and can be rescued by inhibition of ethylene biosynthesis, but not by blocking the perception of ethylene (Xu *et al.*, 2008). These data suggest that cell wall synthesis might be regulated by plasma membrane–cell wall anchors through hormone-signalling pathways, but the mechanism underpinning these connections is unknown.

## Conclusion and future perspectives

The interdependence of the cell wall and intracellular compartments is important for plant development and relies on adequate connections between the cell wall and plasma membrane. Important questions remain to be answered in the fields of secretion and internalization of cell wall material, and of the coordination between cell wall synthesis and degradation. One especially puzzling question relates to how the plasma membrane lipids and CSCs interact during cellulose synthesis. Indeed, the way in which plasma membrane lipids affect protein function *in vivo*, and how they change during vesicle trafficking, is largely unknown. Interestingly, cell wall architecture influences the activity of the CSCs. Moreover, the motility and localization of other plasma membrane proteins are also controlled by cell wall polymers, probably by physical connections which have not been fully elucidated. Therefore, these plasma membrane proteins and CSCs might act as plasma membrane–cell wall anchors with putative roles in cell adhesion.

Cellular adhesion is essential for maintaining the multicellular structure and for signal transduction in eukaryotes. It is plausible that the plasma membrane proteins involved in cell adhesion act as mechanosensors and as receptors of pathogen effectors (Baluška *et al.*, 2003). It is becoming increasingly clear that plants share a similar mechanism to that described in animals for perceiving apoplastic signals by plasma membrane–cell wall adhesion domains, and integrate them into cytoskeletal and hormonal responses to regulate cell expansion (Baluška *et al.*, 2003; Sampathkumar *et al.*, 2014). While plants have many putative plasma membrane–cell wall anchoring proteins, similar to those identified in cell adhesion in other eukaryotes, much remains to be investigated. For example, which plant proteins contain RGD motifs? And do these RGD-containing proteins bind to cell wall polysaccharides? Moreover, little information is available about the downstream effects, and effectors, of changes in cell adhesion properties, e.g. impacts on the cytoskeleton and hormone responses. In addition, a complete characterization of plasma membrane–cell wall adhesion needs a multicellular perspective, i.e. including influences from neighbouring cells. We envision that we will get closer to such insights by the development and combination of proteomic, biochemical, biomechanical, and imaging tools. Thus, we expect that a deeper understanding of the plant plasma membrane–cell wall connection will increase our knowledge about how plant tissues are organized through cell adhesion and cell–cell communication



## Acknowledgements

We apologize to those colleagues whose work we could not cover due to space limitations. This work was supported by Chinese Scholarship Council (ZL), the Max Planck Gesellschaft (SP), and the Deutsche Forschungsgemeinschaft (CSR; PE1642/5-1).

## References

- Alassimone J, Naseer S, Geldner N.** 2010. A developmental framework for endodermal differentiation and polarity. *Proceedings of the National Academy of Sciences, USA* **107**, 5214–5219.
- Alexandersson E, Saalbach G, Larsson C, Kjellbom P.** 2004. *Arabidopsis* plasma membrane proteomics identifies components of transport, signal transduction and membrane trafficking. *Plant and Cell Physiology* **45**, 1543–1556.
- Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P, eds.** 2002. *Molecular biology of the cell*. Ed 4. New York: Garland Science.
- Alejandro S, Lee Y, Tohge T, et al.** 2012. AtABCG29 is a monolignol transporter involved in lignin biosynthesis. *Current Biology* **22**, 1207–1212.
- Atmodjo MA, Hao Z, Mohnen D.** 2013. Evolving views of pectin biosynthesis. *Annual Review of Plant Biology* **64**, 747–779.
- Baluška F, Hlavacka A, Šamaj J, Palme K, Robinson DG, Matoh T, McCurdy DW, Menzel D, Volkmann D.** 2002. F-actin-dependent endocytosis of cell wall pectins in meristematic root cells. Insights from brefeldin A-induced compartments. *Plant Physiology* **130**, 422–431.
- Baluška F, Liners F, Hlavacka A, Schlicht M, Van Cutsem P, McCurdy DW, Menzel D.** 2005. Cell wall pectins and xyloglucans are internalized into dividing root cells and accumulate within cell plates during cytokinesis. *Protoplasma* **225**, 141–155.
- Baluška F, Šamaj J, Wojtaszek P, Volkmann D, Menzel D.** 2003. Cytoskeleton-plasma membrane-cell wall continuum in plants. Emerging links revisited. *Plant Physiology* **133**, 482–491.
- Bashline L, Li S, Anderson CT, Lei L, Gu Y.** 2013. The endocytosis of cellulose synthase in *Arabidopsis* is dependent on mu2, a clathrin-mediated endocytosis adaptin. *Plant Physiology* **163**, 150–160.
- Bashline L, Li S, Gu Y.** 2014. The trafficking of the cellulose synthase complex in higher plants. *Annals of Botany*.
- Bessueille L, Sindt N, Guichardant M, Djerbi S, Teeri TT, Bulone V.** 2009. Plasma membrane microdomains from hybrid aspen cells are involved in cell wall polysaccharide biosynthesis. *Biochemical Journal* **420**, 93–103.
- Boerjan W, Ralph J, Baucher M.** 2003. Lignin biosynthesis. *Annual Review of Plant Biology* **54**, 519–546.
- Borner GH, Lilley KS, Stevens TJ, Dupree P.** 2003. Identification of glycosylphosphatidylinositol-anchored proteins in *Arabidopsis*. A proteomic and genomic analysis. *Plant Physiology* **132**, 568–577.
- Borner GH, Sherrier DJ, Stevens TJ, Arkin IT, Dupree P.** 2002. Prediction of glycosylphosphatidylinositol-anchored proteins in *Arabidopsis*. A genomic analysis. *Plant Physiology* **129**, 486–499.
- Borner GH, Sherrier DJ, Weimar T, Michaelson LV, Hawkins ND, Macaskill A, Napier JA, Beale MH, Lilley KS, Dupree P.** 2005. Analysis of detergent-resistant membranes in *Arabidopsis*. Evidence for plasma membrane lipid rafts. *Plant Physiology* **137**, 104–116.
- Bouwmeester K, de Sain M, Weide R, Gouget A, Klamer S, Canut H, Govers F.** 2011. The lectin receptor kinase LecRK-I.9 is a novel *Phytophthora* resistance component and a potential host target for a RXLR effector. *PLoS Pathogens* **7**, e1001327.
- Bringmann M, Li E, Sampathkumar A, Kocabek T, Hauser MT, Persson S.** 2012. POM-POM2/cellulose synthase interacting1 is essential for the functional association of cellulose synthase and microtubules in *Arabidopsis*. *The Plant Cell* **24**, 163–177.
- Brutus A, Sicilia F, Macone A, Cervone F, De Lorenzo G.** 2010. A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. *Proceedings of the National Academy of Sciences, USA* **107**, 9452–9457.
- Canut H, Carrasco A, Galaud J-P, Cassan C, Bouyssou H, Vita N, Ferrara P, Pont-Lezica R.** 1998. High affinity RGD-binding sites at the plasma membrane of *Arabidopsis thaliana* links the cell wall. *The Plant Journal* **16**, 63–71.
- Cao Y, Tang X, Giovannoni J, Xiao F, Liu Y.** 2012. Functional characterization of a tomato *COBRA-like* gene functioning in fruit development and ripening. *BMC Plant Biology* **12**, 211.
- Capron A, Gourgues M, Neiva LS, et al.** 2008. Maternal control of male-gamete delivery in *Arabidopsis* involves a putative GPI-anchored protein encoded by the *LORELEI* gene. *The Plant Cell* **20**, 3038–3049.
- Cheung AY, Wu HM.** 2011. THESEUS 1, FERONIA and relatives: a family of cell wall-sensing receptor kinases? *Current Opinion in Plant Biology* **14**, 632–641.
- Collings DA, Gebbie LK, Howles PA, Hurley UA, Birch RJ, Cork AH, Hocart CH, Arioli T, Williamson RE.** 2008. *Arabidopsis* dynamin-like protein DRP1A: a null mutant with widespread defects in endocytosis, cellulose synthesis, cytokinesis, and cell expansion. *Journal of Experimental Botany* **59**, 361–376.
- Coppinger P, Repetti PP, Day B, Dahlbeck D, Mehlert A, Staskawicz BJ.** 2004. Overexpression of the plasma membrane-localized NDR1 protein results in enhanced bacterial disease resistance in *Arabidopsis thaliana*. *The Plant Journal* **40**, 225–237.
- Cosgrove DJ.** 2005. Growth of the plant cell wall. *Nature Reviews Molecular Cell Biology* **6**, 850–861.
- Cvrčková F.** 2013. Formins and membranes: anchoring cortical actin to the cell wall and beyond. *Frontiers in Plant Science* **4**, 436.
- Dai X, You C, Chen G, Li X, Zhang Q, Wu C.** 2011. *OsBC1L4* encodes a COBRA-like protein that affects cellulose synthesis in rice. *Plant Molecular Biology* **75**, 333–345.
- DeBolt S, Gutierrez R, Ehrhardt DW, Somerville C.** 2007. Nonmotile cellulose synthase subunits repeatedly accumulate within localized regions at the plasma membrane in *Arabidopsis* hypocotyl cells following 2,6-dichlorobenzonitrile treatment. *Plant Physiology* **145**, 334–338.
- Decreux A, Messiaen J.** 2005. Wall-associated kinase WAK1 interacts with cell wall pectins in a calcium-induced conformation. *Plant and Cell Physiology* **46**, 268–278.
- Diotallevi F, Mulder B.** 2007. The cellulose synthase complex: a polymerization driven supramolecular motor. *Biophysical Journal* **92**, 2666–2673.
- Drakakaki G, Dandekar A.** 2013. Protein secretion: how many secretory routes does a plant cell have? *Plant Science* **203–204**, 74–78.
- Driouich A, Follet-Gueye ML, Bernard S, Kousar S, Chevalier L, Vitré-Gibouin M, Lerouxel O.** 2012. Golgi-mediated synthesis and secretion of matrix polysaccharides of the primary cell wall of higher plants. *Frontiers in Plant Science* **3**, 79.
- Elkins T, Hortsch M, Bieber AJ, Snow PM, Goodman CS.** 1990. Drosophila Fasciclin-I is a novel homophilic adhesion molecule that along with Fasciclin-III can mediate cell sorting. *Journal of Cell Biology* **110**, 1825–1832.
- Ellis M, Egelund J, Schultz CJ, Bacic A.** 2010. Arabinogalactan-proteins: key regulators at the cell surface? *Plant Physiology* **153**, 403–419.
- Fan X, Yang C, Klisch D, Ferguson A, Bhaellero RP, Niu X, Wilson ZA.** 2014. ECHIDNA protein impacts on male fertility in *Arabidopsis* by mediating trans-Golgi network secretory trafficking during anther and pollen development. *Plant Physiology* **164**, 1338–1349.
- Feraru E, Feraru MI, Kleine-Vehn J, Martinieri A, Mouille G, Vanneste S, Vernhettes S, Runions J, Friml J.** 2011. PIN polarity maintenance by the cell wall in *Arabidopsis*. *Current Biology* **21**, 338–343.
- Frigerio L.** 2010. Plant exocytosis, endocytosis and membrane recycling in turgid cells. eLS doi: 10.1002/9780470015902.a0001676.pub2
- Fujikura U, Elsaesser L, Breuninger H, Sanchez-Rodriguez C, Ivakov A, Laux T, Findlay K, Persson S, Lenhard M.** 2014. Atkinesin-13A modulates cell-wall synthesis and cell expansion in *Arabidopsis thaliana* via the THESEUS1 pathway. *PLoS Genetics* **10**, e1004627.
- Geisler DA, Sampathkumar A, Mutwil M, Persson S.** 2008. Laying down the bricks: logistic aspects of cell wall biosynthesis. *Current Opinion in Plant Biology* **11**, 647–652.
- Gendre D, McFarlane HE, Johnson E, Mouille G, Sjodin A, Oh J, Levesque-Tremblay G, Watanabe Y, Samuels L, Bhalarao RP.** 2013. Trans-Golgi network localized ECHIDNA/Ypt interacting protein complex is



required for the secretion of cell wall polysaccharides in *Arabidopsis*. *The Plant Cell* **25**, 2633–2646.

**Gendre D, Oh J, Boutte Y, Best JG, Samuels L, Nilsson R, Uemura T, Marchant A, Bennett MJ, Grebe M, Bhalarao RP.** 2011. Conserved *Arabidopsis* ECHIDNA protein mediates trans-Golgi-network trafficking and cell elongation. *Proceedings of the National Academy of Sciences, USA* **108**, 8048–8053.

**Gillmor CS, Lukowitz W, Brininstool G, Sedbrook JC, Hamann T, Poindexter P, Somerville C.** 2005. Glycosylphosphatidylinositol-anchored proteins are required for cell wall synthesis and morphogenesis in *Arabidopsis*. *The Plant Cell* **17**, 1128–1140.

**Goodman CS, Davis GW, Zito K.** 1997. The many faces of fasciclin II: Genetic analysis reveals multiple roles for a cell adhesion molecule during the generation of neuronal specificity. *Cold Spring Harbor Symposia on Quantitative Biology* **62**, 479–491.

**Gouget A, Senchou V, Govers F, Sanson A, Barre A, Rouge P, Pont-Lezica R, Canut H.** 2006. Lectin receptor kinases participate in protein-protein interactions to mediate plasma membrane-cell wall adhesions in *Arabidopsis*. *Plant Physiology* **140**, 81–90.

**Griffith OH, Ryan M.** 1999. Bacterial phosphatidylinositol-specific phospholipase C: structure, function, and interaction with lipids. *Biochimica et Biophysica Acta* **1441**, 237–254.

**Griffiths JS, Tsai AY, Xue H, Voiniciuc C, Šola K, Seifert GJ, Mansfield SD, Haughn GW.** 2014. SALT-OVERLY SENSITIVE5 mediates *Arabidopsis* seed coat mucilage adherence and organization through pectins. *Plant Physiology* **165**, 991–1004.

**Gutierrez R, Lindeboom JJ, Paredez AR, Emons AM, Ehrhardt DW.** 2009. *Arabidopsis* cortical microtubules position cellulose synthase delivery to the plasma membrane and interact with cellulose synthase trafficking compartments. *Nature Cell Biology* **11**, 797–806.

**Haigler CH, Grimson MJ, Gervais J, Le Moigne N, Höfte H, Monasse B, Navard P.** 2014. Molecular modeling and imaging of initial stages of cellulose fibril assembly: evidence for a disordered intermediate stage. *PLoS ONE* **9**, e93981.

**Harpaz-Saad S, McFarlane HE, Xu S, Divi UK, Forward B, Western TL, Kieber JJ.** 2011. Cellulose synthesis via the FEI2 RLK/SOS5 pathway and cellulose synthase 5 is required for the structure of seed coat mucilage in *Arabidopsis*. *The Plant Journal* **68**, 941–953.

**Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR.** 2014. A peptide hormone and its receptor protein kinase regulate plant cell expansion. *Science* **343**, 408–411.

**Hecht K.** 1912. Studien uiber den Vorgang der Plasmolyse. *Beitrage zur Biologie der Pflanzen II* **37**.

**Hématy K, Sado PE, Van Tuinen A, Rochange S, Desnos T, Balzergue S, Pelletier S, Renou JP, Hofte H.** 2007. A receptor-like kinase mediates the response of *Arabidopsis* cells to the inhibition of cellulose synthesis. *Current Biology* **17**, 922–931.

**Hijazi M, Roujol D, Nguyen-Kim H, Castillo LdRC, Saland E, Jamet E, Albenne C.** 2014. Arabinogalactan protein 31 (AGP31), a putative network-forming protein in *Arabidopsis thaliana* cell walls? *Annals of Botany* **114**, 1087–1097.

**Huang GQ, Gong SY, Xu WL, Li W, Li P, Zhang CJ, Li DD, Zheng Y, Li FG, Li XB.** 2013. A fasciclin-like arabinogalactan protein, GhFLA1, is involved in fiber initiation and elongation of cotton. *Plant Physiology* **161**, 1278–1290.

**Ivakov A, Persson S.** 2013. Plant cell shape: modulators and measurements. *Frontiers in Plant Science* **4**, 439.

**Jamet E, Canut H, Boudart G, Pont-Lezica RF.** 2006. Cell wall proteins: a new insight through proteomics. *Trends in Plant Science* **11**, 33–39.

**Johnson KL, Jones BJ, Bacic A, Schultz CJ.** 2003. The fasciclin-like arabinogalactan proteins of *Arabidopsis*. A multigene family of putative cell adhesion molecules. *Plant Physiology* **133**, 1911–1925.

**Johnson KL, Kibble NA, Bacic A, Schultz CJ.** 2011. A fasciclin-like arabinogalactan-protein (FLA) mutant of *Arabidopsis thaliana*, *fla1*, shows defects in shoot regeneration. *PLoS ONE* **6**, e25154.

**Kang BH, Busse JS, Bednarek SY.** 2003. Members of the *Arabidopsis* dynamin-like gene family, ADL1, are essential for plant cytokinesis and polarized cell growth. *The Plant Cell* **15**, 899–913.

**Kang J, Park J, Choi H, Burla B, Kretschmar T, Lee Y, Martinoia E.** 2011. Plant ABC transporters. *The Arabidopsis Book* **9**, e0153.

**Kawamoto T, Noshiro M, Shen M, Nakamasu K, Hashimoto K, Kawashima-Ohya Y, Gotoh O, Kato Y.** 1998. Structural and phylogenetic analyses of RGD-CAP/beta ig-h3, a fasciclin-like adhesion protein expressed in chick chondrocytes. *Biochimica et Biophysica Acta* **1395**, 288–292.

**Ketelaar T, Galway ME, Mulder BM, Emons AM.** 2008. Rates of exocytosis and endocytosis in *Arabidopsis* root hairs and pollen tubes. *Journal of Microscopy* **231**, 265–273.

**Kim SY, Xu ZY, Song K, Kim DH, Kang H, Reichardt I, Sohn EJ, Friml J, Juergens G, Hwang I.** 2013. Adaptor protein complex 2-mediated endocytosis is crucial for male reproductive organ development in *Arabidopsis*. *The Plant Cell* **25**, 2970–2985.

**Knepper C, Savory EA, Day B.** 2011a. *Arabidopsis* NDR1 is an integrin-like protein with a role in fluid loss and plasma membrane-cell wall adhesion. *Plant Physiology* **156**, 286–300.

**Knepper C, Savory EA, Day B.** 2011b. The role of NDR1 in pathogen perception and plant defense signaling. *Plant Signal and Behavior* **6**, 1114–1116.

**Kohorn BD, Kohorn SL.** 2012. The cell wall-associated kinases, WAKs, as pectin receptors. *Frontiers in Plant science* **3**, 88.

**Kohorn BD, Kohorn SL, Saba NJ, Martinez VM.** 2014. Requirement for pectin methyl esterase and preference for fragmented over native pectins for wall-associated kinase-activated, EDS1/PAD4-dependent stress response in *Arabidopsis*. *The Journal of Biological Chemistry* **289**, 18978–18986.

**Kulich I, Cole R, Drdova E, Cvřicková F, Soukup A, Fowler J, Žárský V.** 2010. *Arabidopsis* exocyst subunits SEC8 and EXO70A1 and exocyst interactor ROH1 are involved in the localized deposition of seed coat pectin. *New Phytologist* **188**, 615–625.

**Lakhan SE, Sabharanjak S, De A.** 2009. Endocytosis of glycosylphosphatidylinositol-anchored proteins. *Journal of Biomedical Science* **16**, 93.

**Lalanne E, Honys D, Johnson A, Borner GH, Lilley KS, Dupree P, Grossniklaus U, Twell D.** 2004. SETH1 and SETH2, two components of the glycosylphosphatidylinositol anchor biosynthetic pathway, are required for pollen germination and tube growth in *Arabidopsis*. *The Plant Cell* **16**, 229–240.

**Lane DR, Wiedemeier A, Peng L, et al.** 2001. Temperature-sensitive alleles of RSW2 link the KORRIGAN endo-1,4-beta-glucanase to cellulose synthesis and cytokinesis in *Arabidopsis*. *Plant Physiology* **126**, 278–288.

**Lang I, Barton DA, Overall RL.** 2004. Membrane-wall attachments in plasmolysed plant cells. *Protoplasma* **224**, 231–243.

**Lefebvre B, Furt F, Hartmann MA, et al.** 2007. Characterization of lipid rafts from *Medicago truncatula* root plasma membranes: a proteomic study reveals the presence of a raft-associated redox system. *Plant Physiology* **144**, 402–418.

**Lei L, Zhang T, Strasser R, et al.** 2014. The *jiaoyao1* mutant is an allele of *korrigan1* that abolishes endoglucanase activity and affects the organization of both cellulose microfibrils and microtubules in *Arabidopsis*. *The Plant Cell* **26**, 2601–2616.

**Lerouxel O, Cavalier DM, Liepman AH, Keegstra K.** 2006. Biosynthesis of plant cell wall polysaccharides - a complex process. *Current Opinion in Plant Biology* **9**, 621–630.

**Li J, Yu M, Geng LL, Zhao J.** 2010. The fasciclin-like arabinogalactan protein gene, FLA3, is involved in microspore development of *Arabidopsis*. *The Plant Journal* **64**, 482–497.

**Li S, Lei L, Somerville CR, Gu Y.** 2012. Cellulose synthase interactive protein 1 (CSI1) links microtubules and cellulose synthase complexes. *Proceedings of the National Academy of Sciences, USA* **109**, 185–190.

**Lindner H, Muller LM, Boisson-Dernier A, Grossniklaus U.** 2012. CrRLK1L receptor-like kinases: not just another brick in the wall. *Current Opinion in Plant Biology* **15**, 659–669.

**Liu C, Mehdy MC.** 2007. A nonclassical arabinogalactan protein gene highly expressed in vascular tissues, AGP31, is transcriptionally repressed by methyl jasmonic acid in *Arabidopsis*. *Plant Physiology* **145**, 863–874.

**Liu L, Shang-Guan K, Zhang B, et al.** 2013. Brittle Culm1, a COBRA-like protein, functions in cellulose assembly through binding cellulose microfibrils. *PLoS Genetics* **9**, e1003704.

**Lü B, Chen F, Gong ZH, Xie H, Liang JS.** 2007a. Integrin-like protein is involved in the osmotic stress-induced abscisic acid biosynthesis in *Arabidopsis thaliana*. *Journal of Integrative Plant Biology* **49**, 540–549.

- Lü B, Chen F, Gong ZH, Xie H, Zhang JH, Liang JS. 2007b. Intracellular localization of integrin-like protein and its roles in osmotic stress-induced abscisic acid biosynthesis in *Zea mays*. *Protoplasma* **232**, 35–43.
- Lü B, Wang J, Zhang Y, Wang H, Liang J, Zhang J. 2012. AT14A mediates the cell wall-plasma membrane-cytoskeleton continuum in *Arabidopsis thaliana* cells. *Journal of Experimental Botany* **63**, 4061–4069.
- MacMillan CP, Mansfield SD, Stachurski ZH, Evans R, Southerton SG. 2010. Fasciclin-like arabinogalactan proteins: specialization for stem biomechanics and cell wall architecture in *Arabidopsis* and *Eucalyptus*. *The Plant Journal* **62**, 689–703.
- Martinière A, Gayral P, Hawes C, Runions J. 2011. Building bridges: formin1 of *Arabidopsis* forms a connection between the cell wall and the actin cytoskeleton. *The Plant Journal* **66**, 354–365.
- Martinière A, Lavagi I, Nageswaran G, *et al.* 2012. Cell wall constrains lateral diffusion of plant plasma-membrane proteins. *Proceedings of the National Academy of Sciences, USA* **109**, 12805–12810.
- Martinière A, Runions J. 2013. Protein diffusion in plant cell plasma membranes: the cell-wall corral. *Frontiers in Plant Science* **4**, 515.
- McFarlane HE, Doring A, Persson S. 2014. The cell biology of cellulose synthesis. *Annual Review of Plant Biology* **65**, 69–94.
- McFarlane HE, Shin JJ, Bird DA, Samuels AL. 2010. *Arabidopsis* ABCG transporters, which are required for export of diverse cuticular lipids, dimerize in different combinations. *The Plant Cell* **22**, 3066–3075.
- Mei Y, Gao HB, Yuan M, Xue HW. 2012. The *Arabidopsis* ARCP protein, CS1, which is required for microtubule stability, is necessary for root and anther development. *The Plant Cell* **24**, 1066–1080.
- Mellersh DG, Heath MC. 2001. Plasma membrane-cell wall adhesion is required for expression of plant defense responses during fungal penetration. *The Plant Cell* **13**, 413–424.
- Miari F, Desprez T, Biot E, Morin H, Belcram K, Hofte H, Gonneau M, Vernhettes S. 2014. Spatio-temporal analysis of cellulose synthesis during cell plate formation in *Arabidopsis*. *The Plant Journal* **77**, 71–84.
- Morel J, Claverol S, Mongrand S, Furt F, Fromentin J, Bessoule JJ, Blein JP, Simon-Plas F. 2006. Proteomics of plant detergent-resistant membranes. *Molecular and Cellular Proteomics* **5**, 1396–1411.
- Morgan JL, Strumillo J, Zimmer J. 2013. Crystallographic snapshot of cellulose synthesis and membrane translocation. *Nature* **493**, 181–186.
- Nagpal P, Quatrano RS. 1999. Isolation and characterization of a cDNA clone from *Arabidopsis thaliana* with partial sequence similarity to integrins. *Gene* **230**, 33–40.
- Nicol F, His I, Jauneau A, Vernhettes S, Canut H, Höfte H. 1998. A plasma membrane-bound putative endo-1,4- $\beta$ -D-glucanase is required for normal wall assembly and cell elongation in *Arabidopsis*. *The EMBO Journal* **17**, 5563–5576.
- Oparka KJ. 1994. Plasmolysis: new insights into an old process. *New Phytologist* **126**, 571–591.
- Owen DM, Williamson D, Rentero C, Gaus K. 2009. Quantitative microscopy: protein dynamics and membrane organisation. *Traffic* **10**, 962–971.
- Paredes AR, Somerville CR, Ehrhardt DW. 2006. Visualization of cellulose synthase demonstrates functional association with microtubules. *Science* **312**, 1491–1495.
- Peng L, Kawagoe Y, Hogan P, Delmer D. 2002. Sitosterol-beta-glucoside as primer for cellulose synthesis in plants. *Science* **295**, 147–150.
- Peskan T, Westermann M, Oelmüller R. 2000. Identification of low-density Triton X-100-insoluble plasma membrane microdomains in higher plants. *European Journal of Biochemistry* **267**, 6989–6995.
- Pont-Lezica RF, McNally JG, Pickard BG. 1993. Wall-to-membrane linkers in onion epidermis- some hypothesis. *Plant, Cell and Environment* **16**, 111–112.
- Proseus TE, Boyer JS. 2005. Turgor pressure moves polysaccharides into growing cell walls of *Chara corallina*. *Annals of Botany* **95**, 967–979.
- Roppolo D, Geldner N. 2012. Membrane and walls: who is master, who is servant? *Current Opinion in Plant Biology* **15**, 608–617.
- Rose JKC. 2003. *The plant cell wall*. Oxford: Blackwell.
- Rosli HG, Zheng Y, Pombo MA, Zhong S, Bombarely A, Fei Z, Collmer A, Martin GB. 2013. Transcriptomics-based screen for genes induced by flagellin and repressed by pathogen effectors identifies a cell wall-associated kinase involved in plant immunity. *Genome Biology* **14**, R139.
- Roudier F, Fernandez AG, Fujita M, *et al.* 2005. COBRA, an *Arabidopsis* extracellular glycosyl-phosphatidyl inositol-anchored protein, specifically controls highly anisotropic expansion through its involvement in cellulose microfibril orientation. *The Plant Cell* **17**, 1749–1763.
- Sampathkumar A, Gutierrez R, McFarlane HE, Bringmann M, Lindeboom J, Emons AM, Samuels L, Ketelaar T, Ehrhardt DW, Persson S. 2013. Patterning and lifetime of plasma membrane-localized cellulose synthase is dependent on actin organization in *Arabidopsis* interphase cells. *Plant Physiology* **162**, 675–688.
- Sampathkumar A, Yan A, Krupinski P, Meyerowitz EM. 2014. Physical forces regulate plant development and morphogenesis. *Current Biology* **24**, R475–R483.
- Samuels AL, Bisalputra T. 1990. Endocytosis in elongating root cells of *Lobelia erinus*. *Journal of Cell Science* **97**, 157–165.
- Sánchez-Rodríguez C, Persson S. 2014. Regulation of cell wall formation by membrane traffic. In: Fukuda H. *Plant cell wall patterning and cell shape*. Hoboken, NJ, USA: John Wiley & Sons, 35–64.
- Schindler M, Meiners S, Cheresch DA. 1989. RGD-dependent linkage between plant cell wall and plasma membrane: consequences for growth. *The Journal of Cell Biology* **108**, 1955–1965.
- Schrick K, Debolt S, Bulone V. 2012. Deciphering the molecular functions of sterols in cellulose biosynthesis. *Frontiers in Plant Science* **3**, 84.
- Schrick K, Fujioka S, Takatsuto S, Stierhof YD, Stransky H, Yoshida S, Jurgens G. 2004. A link between sterol biosynthesis, the cell wall, and cellulose in *Arabidopsis*. *The Plant Journal* **38**, 227–243.
- Sedbrook JC, Carroll KL, Hung KF, Masson PH, Somerville CR. 2002. The *Arabidopsis* SKU5 gene encodes an extracellular glycosyl phosphatidylinositol-anchored glycoprotein involved in directional root growth. *The Plant Cell* **14**, 1635–1648.
- Seifert GJ, Roberts K. 2007. The biology of arabinogalactan proteins. *Annual Review of Plant Biology* **58**, 137–161.
- Seifert GJ, Xue H, Acet T. 2014. The *Arabidopsis thaliana* FASCICLIN LIKE ARABINO GALACTAN PROTEIN 4 gene acts synergistically with abscisic acid signalling to control root growth. *Annals of Botany* **114**, 1125–1133.
- Senchou V, Weide R, Carrasco A, Bouyssou H, Pont-Lezica R, Govers F, Canut H. 2004. High affinity recognition of a *Phytophthora* protein by *Arabidopsis* via an RGD motif. *Cellular and Molecular Life Sciences* **61**, 502–509.
- Sethaphong L, Haigler CH, Kubicki JD, Zimmer J, Bonetta D, DeBolt S, Yingling YG. 2013. Tertiary model of a plant cellulose synthase. *Proceedings of the National Academy of Sciences, USA* **110**, 7512–7517.
- Shi H, Kim Y, Guo Y, Stevenson B, Zhu JK. 2003. The *Arabidopsis* SOS5 locus encodes a putative cell surface adhesion protein and is required for normal cell expansion. *The Plant Cell* **15**, 19–32.
- Shih H-W, Miller ND, Dai C, Spalding EP, Monshausen GB. 2014. The receptor-like kinase FERONIA is required for mechanical signal transduction in *Arabidopsis* seedlings. *Current Biology* **24**, 1887–1892.
- Snow PM, Bieber AJ, Goodman CS. 1989. Fasciclin III: A novel homophilic adhesion molecule in *Drosophila*. *Cell* **59**, 313–323.
- Somerville C, Bauer S, Brininstool G, *et al.* 2004. Toward a systems approach to understanding plant cell walls. *Science* **306**, 2206–2211.
- Sorek N, Sorek H, Kijac A, Szemenyei HJ, Bauer S, Hematy K, Wemmer DE, Somerville CR. 2014. The *Arabidopsis* COBRA protein facilitates cellulose crystallization at the plasma membrane. *The Journal of Biological Chemistry* **289**, 34911–34920.
- Takahashi J, Rudsander UJ, Hedenstrom M, *et al.* 2009. KORRIGAN1 and its aspen homolog PttCel9A1 decrease cellulose crystallinity in *Arabidopsis* stems. *Plant and Cell Physiology* **50**, 1099–1115.
- Tan L, Eberhard S, Pattathil S, *et al.* 2013. An *Arabidopsis* cell wall proteoglycan consists of pectin and arabinoxylan covalently linked to an arabinogalactan protein. *The Plant Cell* **25**, 270–287.
- Tsakamoto T, Qin Y, Huang Y, Dunatunga D, Palanivelu R. 2010. A role for LORELEI, a putative glycosylphosphatidylinositol-anchored protein,

in *Arabidopsis thaliana* double fertilization and early seed development. *The Plant Journal* **62**, 571–588.

**Tsukaya H.** 2006. Mechanism of leaf-shape determination. *Annual Review of Plant Biology* **57**, 477–496.

**Vain T, Crowell EF, Timpano H, et al.** 2014. The cellulase KORRIGAN is part of the cellulose synthase complex. *Plant Physiology* **165**, 1521–1532.

**Van Damme D, Coutuer S, De Rycke R, Bouget FY, Inzé D, Geelen D.** 2006. Somatic cytokinesis and pollen maturation in *Arabidopsis* depend on TPLATE, which has domains similar to coat proteins. *The Plant Cell* **18**, 3502–3518.

**van Hengel AJ, Roberts K.** 2003. AtAGP30, an arabinogalactan-protein in the cell walls of the primary root, plays a role in root regeneration and seed germination. *The Plant Journal* **36**, 256–270.

**Vogel JP, Raab TK, Schiff C, Somerville SC.** 2002. PMR6, a pectate lyase-like gene required for powdery mildew susceptibility in *Arabidopsis*. *The Plant Cell* **14**, 2095–2106.

**Vogel JP, Raab TK, Somerville CR, Somerville SC.** 2004. Mutations in PMR5 result in powdery mildew resistance and altered cell wall composition. *The Plant Journal* **40**, 968–978.

**Wagner TA, Kohorn BD.** 2001. Wall-associated kinases are expressed throughout plant development and are required for cell expansion. *The Plant Cell* **13**, 303–318.

**Wolf S, Hématy K, Höfte H.** 2012a. Growth control and cell wall signaling in plants. *Annual Review of Plant Biology* **63**, 381–407.

**Wolf S, Mravec J, Greiner S, Mouille G, Höfte H.** 2012b. Plant cell wall homeostasis is mediated by brassinosteroid feedback signaling. *Current Biology* **22**, 1732–1737.

**Xie L, Yang C, Wang X.** 2011. Brassinosteroids can regulate cellulose biosynthesis by controlling the expression of *CESA* genes in *Arabidopsis*. *Journal of Experimental Botany* **62**, 4495–4506.

**Xiong G, Li R, Qian Q, Song X, Liu X, Yu Y, Zeng D, Wan J, Li J, Zhou Y.** 2010. The rice dynamin-related protein DRP2B mediates membrane trafficking, and thereby plays a critical role in secondary cell wall cellulose biosynthesis. *The Plant Journal* **64**, 56–70.

**Xiong L, Ishitani M, Lee H, Zhu JK.** 2001. The *Arabidopsis* *LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress-responsive gene expression. *The Plant Cell* **13**, 2063–2083.

**Xiong L, Lee H, Ishitani M, Zhu JK.** 2002. Regulation of osmotic stress-responsive gene expression by the *LOS6/ABA1* locus in *Arabidopsis*. *The Journal of Biological Chemistry* **277**, 8588–8596.

**Xu SL, Rahman A, Baskin TI, Kieber JJ.** 2008. Two leucine-rich repeat receptor kinases mediate signaling, linking cell wall biosynthesis and ACC synthase in *Arabidopsis*. *The Plant Cell* **20**, 3065–3079.

**Yang J, Sardar HS, McGovern KR, Zhang Y, Showalter AM.** 2007. A lysine-rich arabinogalactan protein in *Arabidopsis* is essential for plant growth and development, including cell division and expansion. *The Plant Journal* **49**, 629–640.

**Young RE, McFarlane HE, Hahn MG, Western TL, Haughn GW, Samuels AL.** 2008. Analysis of the Golgi apparatus in *Arabidopsis* seed coat cells during polarized secretion of pectin-rich mucilage. *The Plant Cell* **20**, 1623–1638.

**Zaban B, Maisch J, Nick P.** 2013. Dynamic actin controls polarity induction de novo in protoplasts. *Journal of Integrative Plant Biology* **55**, 142–159.

**Žárský V, Cvřčková F, Martin Potocký M, Hála M.** 2009. Exocytosis and cell polarity in plants - exocyst and recycling domains. *New Phytologist* **183**, 255–272.

**Zauber H, Burgos A, Garapati P, Schulze WX.** 2014. Plasma membrane lipid-protein interactions affect signaling processes in sterol-biosynthesis mutants in *Arabidopsis thaliana*. *Frontiers in Plant Science* **5**, 78.

**Zhang Y, Persson S, Hirst J, Robinson MS, van Damme D, Sanchez-Rodriguez C.** 2014. Change your Tplate, change your fate: plant CME and beyond. *Trends in Plant Science* doi: 10.1016/j.tplants.2014.09.002

**Zhang Y, Yang J, Showalter AM.** 2011. AtAGP18 is localized at the plasma membrane and functions in plant growth and development. *Planta* **233**, 675–683.